

Title: Distressing dreams and risk of Parkinson's disease.

Author: Dr Abidemi I. Otaiku^{1,2} MD

1 Department of Neurology, Birmingham City Hospital, Birmingham, UK

2 Centre for Human Brain Health, University of Birmingham, Birmingham, UK

Correspondence:

Dr Abidemi Otaiku, Department of Neurology, Birmingham City Hospital, Dudley Road,
Birmingham B18 7QH, UK. 0121 554 3801. a.otaiku@nhs.net

Word count: 3,454

Abstract

Background

Parkinson's disease (PD) is associated with alterations to the phenomenology of dreaming - including an increased frequency of distressing dreams. Whether distressing dreams might precede the development of PD is unknown. This study investigated the association between frequent distressing dreams and the risk of incident PD.

Methods

3,830 men from the Osteoporotic Fractures in Men Study (MrOS), a population-based cohort, who were free from PD at baseline (2003-2011) and completed item 5h of the Pittsburgh Sleep Quality Index - which probes the frequency of distressing dreams in the past month, were included in this analysis. Incident PD was based on doctor diagnosis. Multivariate logistic regression was used to estimate odds ratios (OR) for incident PD according to distressing dream frequency, with adjustment for potential confounders.

Findings

During a mean follow-up of 7.3 years, 91 (2.4%) cases of incident PD were identified. Participants with frequent distressing dreams had a 2-fold risk for incident PD (OR, 2.03; 95% CI, 1.1-3.6, $P=0.02$). When stratified by follow-up time, frequent distressing dreams were associated with a 6-fold risk for incident PD during the first 4 years after baseline (OR, 5.60; 95% CI, 2.1-15.1; $P<0.001$), however this association had attenuated during the subsequent 8 years (OR, 1.24; 95% CI, 0.5-3.0; $P=0.64$).

Interpretation

In this prospective cohort, frequent distressing dreams were associated with an increased risk for incident PD. The association was only significant within the 4 years prior to diagnosis, which suggests that frequent distressing dreams may be a prodromal symptom of PD.

Funding

The study received no external funding.

Research in context

Evidence before the study

Previous studies have shown that Parkinson's disease (PD) is associated with alterations to the phenomenology of dreaming – including an increased frequency of distressing dreams (nightmares or bad dreams). Approximately one quarter of PD patients report having frequent nightmares; the majority of whom have them sufficiently frequent to be considered a clinical disorder. Given that frequent distressing dreams are more common in people with PD than in the general population, and are associated with more rapid disease progression in PD; it is plausible that they may also be associated with an increased risk for developing PD.

Added value of this study

In this population-based prospective study of 3,818 older men without PD, it was found that participants with self-reported frequent distressing dreams at baseline (≥ 1 /week), had a 2-fold risk for incident PD during a 12-year follow-up. In the first 4-years after baseline, frequent distressing dreams were associated with a 6-fold risk for incident PD, however this association had attenuated and was no longer significant during the subsequent 8 years.

Implications of all the available evidence

Frequent distressing dreams in older men are associated with an increased risk for developing PD and may represent prodromal symptoms. Screening for late onset distressing dreams may help to identify individuals at risk of developing PD.

Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disease and the fastest growing neurological disorder in the world.¹ With the ageing of the world's population, decline in smoking rates, and changes in environmental risk factors linked to PD, it is projected that the global prevalence of PD could double within the next decade.¹ Given that PD remains incurable, identifying individuals who are at high risk for developing PD - in whom interventions could be targeted, is now considered an urgent priority.

Intriguingly, cross-sectional studies have consistently shown that PD is associated with alterations to the phenomenology of dreaming.² This includes changes to dream content (increased negative emotions, increased physical aggression and higher prevalence of animal characters),³ as well as the development of rapid eye movement sleep behaviour disorder (RBD) – or “dream enactment behaviours” (DEBs).² In addition, people with PD are approximately four times more likely than adults in the general population, to experience nightmares sufficiently frequent to be considered a clinical disorder.⁴

Although recent research has shown that a quarter of patients with PD may experience frequent distressing dreams (nightmares or bad dreams) from the time of diagnosis,^{5,6} and that these patients may be at higher risk for early motor and cognitive decline;⁶ it remains to be investigated whether frequent distressing dreams in the general population may be associated with the risk of developing PD.

This study investigated the hypothesis that frequent distressing dreams would be associated with an increased risk for developing PD, and tested this association using data from the Osteoporotic Fractures in Men (MrOS) study,⁷ with up to 12-years follow-up for incident PD.

Methods

Study sample

MrOS is an observational, longitudinal cohort study that enrolled 5994 community-dwelling men aged 65 years or older at six clinical centres in the United States, including: Birmingham, Minneapolis, Palo Alto, Pittsburgh, Portland, and San Diego.⁹ The baseline assessments were conducted between 2000-2002 (Visit 1).

The present analysis used data from: Sleep Visit 1(2003-2005), Visit 2 (2005-2006), Visit 3 (2007-2009), Interim 2 (2009-2011), Sleep Visit 2 (2009-2012), and Visit 4 (2014-2016). Sleep Visit 1 and Interim 2 were considered as the baseline for this study, since these were the earliest visits where information on distressing dreams and history of PD were simultaneously assessed. Interim 2 was considered baseline for the men that did not participate in Sleep Visit 1.

The analysis was limited to participants who completed the questionnaire item on distressing dreams at baseline, were free from PD, had no missing data for any covariates included in the primary analysis, and took part in at least one follow-up after baseline. This yielded an analytic sample comprising 3,818 men, including 3,006 men from Sleep Visit 1 and 812 men from Interim 2.

Distressing dreams

Participants completed the Pittsburgh Sleep Quality Index (PSQI) at baseline, a validated self-report questionnaire for assessing subjective sleep quality and disturbances.⁸

Distressing dreams were assessed using item 5h of the PSQI: "During the past month, how often have you had trouble sleeping because you have bad dreams?". No definition of bad dreams was provided. The response options included: (0) "not during the past month, (1) "Less than once a week", (2) "Once or twice a week" and (3) "Three or more

times a week" (3). Participants experiencing distressing dreams at least once a week were considered to have frequent distressing dreams, consistent with previous studies.⁹

Ascertainment of incident PD

During the 12-year follow-up, participants were asked at each clinical visit or questionnaire-based contact, to report whether they had ever been diagnosed with PD by a doctor. Follow-up time was calculated for each participant as the time to the most recent collection of data on PD diagnosis.

Covariates

Covariates that might confound the association between distressing dreams and incident PD were chosen based on relevant publications.⁹⁻¹¹ These included: age in years (continuous), ethnicity (white, non-white), educational qualifications (college degree/high school degree/none), smoking status (current, past, never), alcohol intake (drinks/week, <1, 1-13, ≥ 14), current depression (yes/no), history of doctor-diagnosed diabetes mellitus (yes/no), history of doctor-diagnosed hypertension (yes/no), sleep onset insomnia (times/week, 0, <1, 1-2, ≥ 3), sleep maintenance insomnia (times/week, 0, <1, 1-2, ≥ 3), self-rated health (good/excellent, poor/fair), physical activity levels (continuous), and use of medications associated with dream content changes (yes/no). In addition, current anxiety (yes/no), excessive daytime sleepiness (yes/no), global cognitive function (continuous) and severity of sleep-disordered breathing (continuous), were available for the participants that took part in Sleep Visit 1.

Age, ethnicity, educational qualifications, alcohol intake, smoking status, history of doctor diagnosed diabetes mellitus, and history of doctor-diagnosed hypertension, were self-reported at baseline. Depressive symptoms were evaluated using the Geriatric Depression Scale, with scores ≥ 6 indicating clinically significant depression.¹² Sleep onset insomnia and sleep maintenance insomnia were assessed using Item 5a and 5b of the

PSQI respectively (scores ranging from 1 "not during the past month", to 4 "three or more times a week").⁸ Level of physical activity was estimated using the Physical Activity Scale for the Elderly (scores range from 0-793, with higher scores indicating greater physical activity levels).¹³ Anxiety symptoms were evaluated using the Goldberg Anxiety Scale, with scores ≥ 5 indicating clinically significant anxiety.¹⁴ Global cognitive function was measured using The Modified Mini-Mental State Examination (scores range from 0-100, with higher scores indicating better cognitive function).¹⁵ Daytime sleepiness was assessed using The Epworth Sleepiness Scale, with scores ≥ 11 indicating excessive daytime sleepiness.¹⁶ Sleep-disordered breathing was assessed using polysomnography and defined with the apnoea-hypopnea index (AHI, apnoeas or hypopneas per hour of sleep with a desaturation of $\geq 3\%$) as previously described.¹⁷ At each visit, participants were asked to bring in all the medications they had used in the last 30 days. Medication use was defined as: antidepressants, benzodiazepines, beta blockers, alpha blockers, antipsychotics, amphetamines, cholinesterase inhibitors, melatonin and nonbenzodiazepine, nonbarbiturate sedative hypnotics.^{2,18}

Statistical analysis

Characteristics of the participants at baseline were compared using χ^2 tests for categorical variables, independent samples t-tests for normally distributed continuous variables, and Mann–Whitney U tests for nonnormally distributed variables.

Multivariate logistic regression was used to obtain odds ratios (RRs) and 95% confidence intervals (CIs) to assess the association between frequent distressing dreams and risk of PD with adjustment for potential confounders. Model 1 was minimally adjusted for age and clinic site. Model 2 additionally adjusted for race, education, smoking status, alcohol intake, depression, sleep onset insomnia, sleep maintenance insomnia, diabetes, hypertension, physical activity levels and self-reported health. Model 3 further adjusted for medication use.

Additional analyses

To explore the temporal relationship between distressing dreams and PD, a stratified analysis was carried using different time-periods: baseline to ~4 years, and 4 to 12 years. As a sensitivity analysis, the primary analysis was repeated after further adjusting for all baseline characteristics that differed between the participants with and without frequent distressing dreams, that were not included as covariates in the primary analysis. These analyses used data from the participants taking part in Sleep Visit 1 (n = 3,006).

Statistical testing was performed two-sided at $P < 0.05$. All analyses were performed using SPSS version 28 (IBM Corp., Armonk, NY).

Ethical considerations

All men provided written informed consent. Each individual site received institutional review board approval before commencement of the study. The present study received approval from the University of Birmingham (Ref No ERN_21-0427).

Role of the funding source

There was no funding source for this study.

Findings

Among 3,818 participants included in this analysis, 368 (9.6%) reported frequent distressing dreams at baseline. Baseline characteristics of the participants stratified by distressing dream frequency are presented in Table 1. The participants with frequent distressing dreams were more likely to be of non-Caucasian ethnicity, have lower education, worse cognitive function, worse physical and mental health, drank less

alcohol, were less physically active, had a higher AHI and were more likely to use medications that can affect dreaming.

During a mean follow-up of 7.3 years, 91 cases of incident PD (2.4%) were identified. Baseline characteristics of the participants stratified by incident PD status are presented in Table 2. At baseline, the incident PD group were more likely to be depressed, had less frequent sleep maintenance insomnia, and had a lower AHI. There were no other differences in characteristics between the participants that did and did not develop PD.

In the fully adjusted logistic regression model (Table 3), compared with participants without frequent distressing dreams at baseline, participants with frequent distressing dreams had a 2-fold risk for incident PD (OR, 2.03; 95% CI, 1.1-3.6, $P = 0.02$).

In the stratified analysis which divided the follow-up into two time-periods (Table 3), frequent distressing dreams were associated with a 6-fold risk for incident PD in the first 4 years after baseline (OR, 5.60; 95% CI, 2.1-15.1; $P < 0.001$). However, this association had attenuated and was no longer significant for PD cases that developed during the subsequent 8 years (OR, 1.24; 95% CI, 0.5-3.0; $P = 0.64$).

In the sensitivity analysis that adjusted for all characteristics that differed between the participants with and without frequent distressing dreams at baseline (Table 3), the association between frequent distressing dreams and incident PD was similar and remained significant (OR, 2.34; 95% CI, 1.2-4.5; $P = 0.01$).

Discussion

In this large prospective study of community-dwelling older men without PD, frequent distressing dreams were associated with an increased risk for incident PD during a 12-year follow-up, accounting for a wide range of possible confounders.

This is the first study to investigate the association between dream content and risk of PD. Of note, a recent population-based prospective study investigated the association between DEBs (the cardinal symptom of idiopathic RBD [iRBD]) and PD risk, and found that DEBs were associated with a 2.9-fold risk for incident PD during a 3-year follow-up.¹⁹ Given that iRBD is considered to be the strongest prodromal marker for PD,²⁰ the finding of a 5.6-fold risk for incident PD among participants with frequent distressing dreams in this study - during an even longer follow-up duration, is a highly significant finding.

The findings of this study are consistent with prior cross-sectional studies, which demonstrated that distressing dreams are very common in people with PD,^{2,3} and that people with PD are significantly more likely to report having frequent distressing dreams than adults in the general population.^{4,5} Indeed, studies have shown that 17% of patients with non-demented PD report weekly nightmares,⁴ and this increases to 78% for patients with PD dementia (PDD).²¹ In contrast, estimates for the prevalence of weekly nightmares in the general population range from 2% to 5%.⁹⁻¹¹

An earlier study found that 80% of patients with PD who reported having frequent intense, vivid and frightening dreams, shortly after the time of diagnosis, believed that the changes to their dream content preceded the onset of their motor symptoms by several years, or even decades.⁵ Thus, the findings of the current study provide prospective evidence to support the patients retrospective recall of prodromal distressing dreams. Given that patients with distressing dreams at the time of diagnosis may be five times more-likely to progress from mild PD to moderate-severe PD (Hoehn & Yahr stage ≥ 3) within 5 years,⁶ and may experience cognitive decline at a rate 32-times faster than patients without dream content changes,⁶ it is probable that a significant proportion of the participants with distressing dreams in this cohort, who later developed PD, would have gone on to experience rapid disease progression in the years following their PD diagnosis.

In this study, frequent distressing dreams were associated with a 6-fold risk for PD during the first 4-years after baseline, though this association had substantially attenuated during the 4-to-12-year time-period. This suggests that late onset distressing dreams, rather than life-long distressing dreams, may be linked with increased PD risk. Interestingly, population-based cross-sectional studies have consistently shown that nightmares in adulthood are most prevalent from around 70 years of age,^{10,11} which is strikingly similar to the median age at which PD symptoms first develop.²² Moreover, between adolescence and middle adulthood, women are significantly more likely than men to experience weekly nightmares, yet from age 65 onwards, there is no longer a gender gap in nightmares.^{10,11} This suggests that the age-related increase in dysphoric dreams is most pronounced in men. Thus, these findings indicate that new onset distressing dreams in older adults, particularly older men, may represent early signs of neurodegeneration in some individuals. This hypothesis is in keeping with the findings from the present study, and is consistent with prior studies, which showed that men with PD have significantly more disturbing dreams than women with PD.³

The neural correlates of dreaming have only recently begun to be identified.²³ There is accumulating evidence that the same brain regions involved in regulating emotions during wakefulness, are also involved in regulating emotions during dreaming.^{24,25} During rapid eye movement (REM) sleep – the sleep stage where nightmares and bad dreams typically occur – activity and blood flow in limbic and paralimbic areas is increased,²⁶ which could explain the characteristic hyperemotionality of dreams.² In contrast, activity in frontoparietal associative networks is decreased during REM sleep,²⁶ which leads to diminished insight and reduced cognitive control.²

A recent study in patients with PD identified that structural changes in the right frontal lobe (right superior frontal gyrus and right anterior cingulate) were correlated with an increased frequency of distressing dreams.²⁷ An earlier study identified that patients with left-onset PD (indicating greater right hemisphere pathology) were more likely to have frequent distressing dreams than patients with right-onset PD (indicating greater left

hemisphere pathology), and also healthy controls; with the latter two groups showing no difference.²⁸ Distressing dreams have also been shown to correlate with poor cognitive function in PD,²⁹ particularly frontal executive function,²⁹ and also with severity of depression and anxiety.³⁰ Given that cognitive impairment and depression usually co-occur in PD, and are each associated with increased PD risk,²⁰ it is plausible that distressing dreams and depressive symptoms in the prodromal stage of PD, are both caused by neurodegeneration of right frontal brain regions, that are involved in the downregulation of negative emotions across conscious states.^{24,25} This hypothesis would be consistent with the neurocognitive model of nightmares.³¹

However, whilst the participants with frequent distressing dreams in this study were indeed more likely to be depressed and to have worse cognitive function (Table 1) - as this hypothesis would predict; after adjusting for these characteristics as covariates, distressing dreams remained significantly associated with incident PD. This suggests that distressing dreams may be a prodromal symptom of PD even in individuals with normal cognition, and who are not depressed. Although, a recent study identified that newly diagnosed patients with PD who reported having recurrent dreams with “aggressive” content (the most frequently reported theme in nightmares),³² did not significantly differ at baseline from the patients without frequent aggressive dreams, with respects to depression or cognitive function.⁶ However, after 5 years, the patients who reported frequent aggressive dreams at baseline, were nearly three times as likely to have developed PDD or mild cognitive impairment, and were both more depressed and more anxious.⁶ Thus, it is possible that distressing dreams may reveal subtle frontal executive deficits, that are not yet detectable by neuropsychological tests or daytime mood disturbances, but may manifest initially as an impaired ability to downregulate negative emotions during REM sleep dreaming.

Strengths and limitations

This study has several strengths, including the prospective design, long follow-up period, use of a well validated questionnaire for assessing habitual sleep disturbances, and inclusion of a wide range of potential confounders. Furthermore, the participants were not selected based on distressing dream frequency or PD diagnosis. Several limitations also warrant discussion. Following previous published studies,³³ this study relied on self-reported doctor diagnosis to determine incident PD and therefore may have missed or misclassified some cases. This misclassification could have led to an underestimation of the association. Second, the questionnaire item used to assess distressing dreams does not clearly distinguish between frequent bad dreams (i.e., distressing dreams without awakenings) and frequent nightmares (i.e., distressing dreams with awakenings). As such, it is not possible to determine whether the associations with PD may vary by distressing dream subtype. Finally, the findings from this study might not be generalisable to women and younger adults.

Conclusions

This study provides evidence for the first time that frequent distressing dreams in community-dwelling older adults, may be associated with an increased risk for developing PD. Participants with frequent distressing dreams were nearly 5 times more likely to develop PD within the initial 4 years of follow-up, however this association had attenuated and was no longer significant during the subsequent 8 years. This suggests that frequent distressing dreams may be a prodromal symptom of PD. As such, screening for late onset distressing dreams in the general population, may help to identify individuals at increased risk of developing a rapidly progressing subtype of PD,⁶ in whom early interventions could be targeted.

Author contributions:

1. Research Project: Conception, Organization, Execution, 2. Statistical Analysis: Design, Execution, 3. Manuscript Preparation

AIO: 1,2,3.

Declaration of Competing Interests:

The author is funded by a National Institute for Health Research academic clinical fellowship.

Data sharing statement

Data from MrOS are available at <https://mrosonline.ucsf.edu>. The analysis dataset for this specific manuscript are also available from the corresponding author upon request.

Funding:

The study received no external funding.

Acknowledgements:

The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health funding. The following institutes provide support: the National Institute on Aging (NIA), the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Center for Advancing Translational Sciences (NCATS), and NIH Roadmap for Medical Research under the following grant numbers: U01 AG027810, U01 AG042124, U01 AG042139, U01 AG042140, U01 AG042143, U01 AG042145, U01 AG042168, U01 AR066160, R01 AG066671, and UL1 TR000128. The National Heart, Lung, and Blood Institute (NHLBI) provides funding for the MrOS Sleep ancillary study "Outcomes of

Sleep Disorders in Older Men" under the following grant numbers: R01 HL071194, R01 HL070848, R01 HL070847, R01 HL070842, R01 HL070841, R01 HL070837, R01 HL070838, and R01 HL070839.

Preprint not peer reviewed

References

1. Dorsey ER, Sherer T, Okun MS, Bloem BR. The Emerging Evidence of the Parkinson Pandemic. *J Parkinsons Dis* 2018;8:S3-S8.
2. Siclari F, Valli K, Arnulf I. Dreams and nightmares in healthy adults and in patients with sleep and neurological disorders. *Lancet Neurol* 2020;10:849-859.
3. Borek LL, Kohn R, Friedman JH. Phenomenology of dreams in Parkinson's disease. *Mov Disord* 2007;22:198-202.
4. Ylikoski A, Martikainen K, Partinen M. Parasomnias and isolated sleep symptoms in Parkinson's disease: a questionnaire study on 661 patients. *J Neurol Sci* 2014;346:204-8.
5. Pont-Sunyer C, Hotter A, Gaig C, et al. The onset of nonmotor symptoms in Parkinson's disease (the ONSET PD study). *Mov Disord* 2015;30:229-37.
6. Otaiku AI. Dream Content Predicts Motor and Cognitive Decline in Parkinson's Disease. *Mov Disord Clin Pract* 2021;8:1041-1051.
7. Orwoll E, Blank JB, Barrett-et al. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study--a large observational study of the determinants of fracture in older men. *Contemp Clin Trials* 2005;26:569-85.
8. Buysse DJ, Reynolds CF 3rd, Monk TH, et al. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.

9. Li SX, Zhang B, Li AM, Wing YK. Prevalence and correlates of frequent nightmares: a community-based 2-phase study. *Sleep* 2010;33:774-80
10. Park D, Kim S, Shin C, et al. Prevalence of and factors associated with nightmares in the elderly in a population based cohort study. *Sleep Med* 2021;78:15-23.
11. Sandman N, Valli K, Kronholm E, Revonsuo A, Laatikainen T, Paunio T. Nightmares: risk factors among the Finnish general adult population. *Sleep* 2015;38:507-14.
12. Sheikh, JI, Yesavage, JA. Geriatric Depression Scale (GDS): recent evidence and development of a shorter version. *Clin Gerontol* 1986;5:165-173.
13. Washburn RA, Smith KW, Jette AM, et al. The Physical Activity Scale for the Elderly (PASE): development and evaluation. *J Clin Epidemiol* 1993;46:153-62.
14. Goldberg D, Bridges K, Duncan-Jones P, et al. Detecting anxiety and depression in general medical settings. *BMJ* 1988;297:897-9.
15. Teng EL, Chui HC. The Modified Mini-Mental State (3MS) examination. *J Clin Psychiatry* 1987;48:314-8.
16. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;14:540-5.
17. Stone KL, Blackwell TL, Ancoli-Israel S, et al. Sleep Disordered Breathing and Risk of Stroke in Older Community-Dwelling Men. *Sleep* 2016;39:531-40.
18. Thompson DF, Pierce DR. Drug-induced nightmares. *Ann Pharmacother* 1999;33:93-8.

19. Yao CW, Saha-Chaudhuri P, Zolfaghari S, Pelletier A, Postuma RB. Phenoconversion from Possible REM Sleep Behavior to Parkinsonism in the Population-Based CLSA. *Mov Disord*. 2021 Jun;36(6):1466-1467.
20. Heinzel S, Berg D, Gasser T, Chen H, Yao C, Postuma RB; MDS Task Force on the Definition of Parkinson's Disease. Update of the MDS research criteria for prodromal Parkinson's disease. *Mov Disord* 2019;34:1464-1470.
21. Pistacchi M, Gioulis M, Contin F, Sanson F, Marsala SZ. Sleep disturbance and cognitive disorder: epidemiological analysis in a cohort of 263 patients. *Neurol Sci* 2014;35:1955-62
22. Breen DP, Evans JR, Farrell K, Brayne C, Barker RA. Determinants of delayed diagnosis in Parkinson's disease. *J Neurol* 2013;260:1978-81.
23. Siclari F, Baird B, Perogamvros L, et al The neural correlates of dreaming. *Nat Neurosci* 2017;20:872-878.
24. Sterpenich V, Perogamvros L, Tononi G, Schwartz S. Fear in dreams and in wakefulness: Evidence for day/night affective homeostasis. *Hum Brain Mapp* 2020;41:840-850.
25. Sikka P, Revonsuo A, Noreika V, Valli K. EEG Frontal Alpha Asymmetry and Dream Affect: Alpha Oscillations over the Right Frontal Cortex during REM Sleep and Presleep Wakefulness Predict Anger in REM Sleep Dreams. *J Neurosci* 2019;39:4775-4784.
26. Maquet P, Péters J, Aerts J, Delfiore G, Degueldre C, Luxen A, Franck G. Functional neuroanatomy of human rapid-eye-movement sleep and dreaming. *Nature* 1996;383:163-6.

27. Radziunas A, Deltuva VP, Tamasauskas A, Gleizniene R, Pranckeviciene A, Petrikonis K, Bunevicius A. Brain MRI morphometric analysis in Parkinson's disease patients with sleep disturbances. *BMC Neurol* 2018;18:88.
28. Stavitsky K, McNamara P, Durso R, Harris E, Auerbach S, Cronin-Golomb A. Hallucinations, dreaming, and frequent dozing in Parkinson disease: impact of right-hemisphere neural networks. *Cogn Behav Neurol* 2008;21:143-9.
29. Bugalho P, Paiva T. Dream features in the early stages of Parkinson's disease. *J Neural Transm* 2011;118:1613-9.
30. Borek LL, Kohn R, Friedman JH. Mood and sleep in Parkinson's disease. *J Clin Psychiatry* 2006;67:958-63.
31. Nielsen T, Levin R. Nightmares: a new neurocognitive model. *Sleep Med Rev* 2007;11:295–310
32. Robert G, Zadra A. Thematic and content analysis of idiopathic nightmares and bad dreams. *Sleep* 2014;37:409-17.
33. Leng Y, Blackwell T, Cawthon PM, Ancoli-Israel S, Stone KL, Yaffe K. Association of Circadian Abnormalities in Older Adults With an Increased Risk of Developing Parkinson Disease. *JAMA Neurol* 2020;77:1270-1278

Table 1. Baseline characteristics by frequency of distressing dreams (DD).

Characteristic	Non-frequent DD (n = 3,450)	Frequent DD (n= 368)	<i>P</i> value
Age (years)	77.0 ± 5.5	77.0 ± 5.1	0.89
White, n (%)	3151 (91.3)	324 (88.0)	0.04
Education, n (%)			0.002
College degree	1978 (57.3)	193 (52.4)	
High school degree	1311 (38.0)	143 (38.9)	
No qualifications	161 (4.7)	32 (8.7)	
Smoking status, n (%)			0.71
Current	66 (1.9)	8 (2.2)	
Past	2008 (58.2)	221 (60.1)	
Never	1376 (39.9)	139 (37.8)	
Alcohol intake, drinks/wk, n (%)			0.02
<1	1606 (46.6)	196 (53.3)	
1-13	1643 (47.6)	160 (43.5)	
≥14	201 (5.8)	12 (3.3)	
Diabetes, n (%)	463 (13.4)	66 (17.9)	0.02
Hypertension, n (%)	1683 (48.8)	225 (61.1)	<.001
Self-reported health, n (% good/excellent)	3064 (88.8)	302 (82.1)	<.001
Physical activity score ^a	145.0 ± 72.0	136.5 ± 74.3	0.01
Depression, n (%) ^b	174 (5.0)	47 (12.8)	<.001
Anxiety, n (%) ^c	206 (7.6)	54 (18.1)	<.001
Sleep onset insomnia, times/wk, n (%)			<.001
0	1822 (52.8)	140 (38.0)	
<1	928 (26.9)	82 (22.3)	
1-2	396 (11.5)	81 (22.0)	
≥3	304 (8.8)	65 (17.7)	
Sleep maintenance insomnia, times/wk, n (%)			<.001

0	493 (14.3)	25 (6.8)	
<1	404 (11.7)	22 (6.0)	
1-2	556 (16.1)	44 (12.0)	
≥3	1997 (57.9)	277 (75.3)	
Excessive daytime sleepiness, n (%) ^d	319 (11.8)	57 (19.1)	<.001
Global cognitive function score ^e	93.0 ± 5.9	90.7 ± 8.4	<.001
Apnoea-hypopnea index, events/hr	18.0 ± 15.8	20.3 ± 17.0	0.04
Medication use, n (%)	1616 (46.8)	224 (60.9)	<.001

Plus-minus values are means ± SD

a Physical Activity Scale for the Elderly score. Scores range from 0-793, with higher scores indicating higher physical activity levels.

b Geriatric Depression Scale ≥6.

c Goldberg Anxiety Scale ≥5. *n* = 2703/298.

d Epworth Sleepiness Scale ≥11. *n* = 2707/299.

e Modified Mini-Mental State Examination score. Scores range from 0-100, with higher scores indicate better cognitive function. *n* = 2705/299

f Measured as number of apnoeas and hypopneas per hour of sleep with a desaturation ≥3 %. *n* = 2527/276.

Table 2. Baseline characteristics by incident Parkinson's disease (PD) status.

Characteristic	No PD (n = 3,727)	Incident PD (n = 91)	<i>P</i> value
Age (years)	77.0 ± 5.5	76.7 ± 5.4	0.51
White, n (%)	3390 (91.0)	85 (93.4)	0.42
Education, n (%)			0.66
College degree	2115 (56.7)	56 (61.5)	
High school degree	1424 (38.2)	30 (33.0)	
No qualifications	188 (5.0)	5 (5.5)	
Smoking status, n (%)			0.60
Current	72 (1.9)	2 (2.2)	
Past	2180 (58.5)	49 (53.8)	
Never	1475 (39.6)	40 (44.0)	
Alcohol intake, drinks/wk, n (%)			0.66
<1	1762 (47.3)	40 (44.0)	
1-13	1756 (47.1)	47 (51.6)	
≥14	209 (5.6)	4 (4.4)	
Diabetes, n (%)	519 (13.9)	10 (11.0)	0.42
Hypertension, n (%)	1858 (49.9)	50 (54.9)	0.34
Self-reported health, n (% good/excellent)	3285 (88.1)	81 (89.0)	0.80
Physical activity score ^a	143.9 ± 71.9	154.0 ± 144.2	0.44
Depression, n (%) ^b	208 (5.6)	13 (14.3)	<.001
Anxiety, n (%) ^c	252 (8.6)	8 (11.0)	0.48
Sleep onset insomnia, times/wk, n (%)			0.28
0	1909 (51.2)	53 (58.2)	
<1	994 (26.7)	16 (17.6)	
1-2	465 (12.5)	12 (13.2)	
≥3	359 (9.6)	10 (11.0)	
Sleep maintenance insomnia, times/wk, n (%)			0.04

0	500 (13.4)	18 (19.8)	
<1	419 (11.2)	7 (7.7)	
1-2	579 (15.5)	21 (23.1)	
≥3	2229 (59.8)	45 (49.5)	
Excessive daytime sleepiness, n (%) ^d	362 (12.4)	14 (18.7)	0.10
Global cognitive function score, n ^e	92.8 ± 6.3	92.4 ± 4.5	0.08
Apnoea-hypopnea index, events/hr ^f	18.4 ± 16.0	13.7 ± 14.3	0.01
Medication use, n (%)	1793 (48.1)	47 (51.6)	0.50

Plus-minus values are means ± SD

a Physical Activity Scale for the Elderly score. Scores range from 0-793, with higher scores indicating higher physical activity levels

b Geriatric Depression Scale ≥6.

c Goldberg Anxiety Scale ≥5. *n* = 2928/73.

d Epworth Sleepiness Scale ≥11. *n* = 2931/75.

e Modified Mini-Mental State Examination score. Scores range from 0-100, with higher scores indicate better cognitive function. *n* = 2929/75.

f Measured as number of apnoeas and hypopneas per hour of sleep with a desaturation ≥3 %. *n* = 2733/70.

Table 3: Frequent distressing dreams and odds ratios (95% CI) for incident Parkinson's disease.

Model	Cases/ <i>n</i>	OR (95%CI)	<i>P</i> value
Main analysis			
Model 1	91/3818	2.02 (1.2, 3.5)	0.01
Model 2	91/3818	2.05 (1.1, 3.6)	0.02
Model 3	91/3818	2.03 (1.1, 3.6)	0.02
Stratified analysis ^a			
Baseline to ~4 years	24/3006	5.60 (2.1, 15.1)	<0.001
4 to 12 years	51/2460	1.24 (0.5, 3.0)	0.64
Sensitivity analysis ^b			
	68/2796	2.32 (1.2, 4.5)	0.01

Model 1: age and clinic site

Model 2: model 1 + race, education, smoking status, alcohol intake, depression, sleep onset insomnia, sleep maintenance insomnia, diabetes, hypertension, physical activity and self-reported health.

Model 3: model 2 + medication use

^a Stratified analysis included participants from Sleep Visit 1 and was adjusted for model 3 covariates.

^b Sensitivity analysis included participants from Sleep Visit 1 and further adjusted for anxiety, cognitive function, excessive daytime sleepiness and apnoea-hypopnea index, in addition to model 3 covariates.